

Symmetric representation of the elements of finite groups

Z. Hasan and
A. Kasouha

*Department of Mathematics, California State University,
5500 University Parkway, San Bernardino, California 92407*

February 2, 2008

Abstract

Many finite groups, including all finite non-abelian simple groups, can be symmetrically generated by involutions. In this paper we give an algorithm to symmetrically represent elements of finite groups and to transform symmetrically represented elements in terms of their permutation representation. In particular, we represent elements of the group $U_3(3) : 2$, where $U_3(3)$ is the projective special unitary group of 3×3 matrices with entries in the field $\mathbb{F}_3$, as a permutation on fourteen letters from the projective general linear group $PGL_2(7)$, followed by a word of length at most two in the fourteen involutory symmetric generators.

1 Introduction

The two general methods for working with groups, permutations and matrices, are inconvenient or unmanageable for large finite groups and in particular for the larger sporadic groups, for example for the Monster group, the least degree of a matrix representation is 196881 and it takes n^3 operations to multiply two matrices of dimension n and the least degree of any permutation representation is 10^{20} . Matrix multiplication for matrices of large size is very time-consuming and, although MAGMA and GAP handle permutations of reasonably large size quite efficiently, recording and transmitting elements is inconvenient. The main purpose of this paper is to give an alternative and more efficient method for working with groups. Double coset enumeration can be performed on groups that possess generating sets of involutions. Curtis has constructed several sporadic groups by manual double coset enumeration; for references see [7] and [8]. Now any finite group generated by a conjugacy class of involutions, and hence all finite non-abelian simple groups have symmetric generating sets of involutions (see [1]). It is this technique of double coset enumeration that allows us to write elements in a much more concise manner. In the first author's work

with Curtis (see [8]), every element of J_1 , usually written as a permutation on 266 letters, was written as an element of $L_2(11)$ followed by a word of length at most 4 in the symmetric generators and a program was written to manipulate elements of J_1 written in this short form.

2 Symmetric generation of a group

Let $\mathcal{G}$ be a group and

$$\mathcal{T} = \{t_1, t_2, \dots, t_n\} \subseteq \mathcal{G},$$

then we define

$$\overline{\mathcal{T}} = \{T_1, T_2, \dots, T_n\},$$

where $T_i = \langle t_i \rangle$, the cyclic subgroup generated by t_i ; we further define $\mathcal{N} = \mathcal{N}_{\mathcal{G}}(\overline{\mathcal{T}})$, the set normalizer in $\mathcal{G}$ of $\overline{\mathcal{T}}$.

If the following two conditions hold:

- (i) $\mathcal{G} = \langle \mathcal{T} \rangle$, and
- (ii) $\mathcal{N}$ permutes $\overline{\mathcal{T}}$ transitively, not necessarily faithfully,

then, following Curtis and Hasan [8], we say that $\mathcal{T}$ is a *symmetric generating set* for $\mathcal{G}$. In these circumstances we call $\mathcal{N}$ the *control subgroup*. Note that (i) and (ii) imply that $\mathcal{G}$ is a homomorphic image of the (infinite) *progenitor*

$$m^{*n} : \mathcal{N},$$

where m^{*n} represents a free product of n copies of the cyclic group C_m , m being the order of t_i , and $\mathcal{N}$ is a group of automorphisms of m^{*n} which permutes the n cyclic subgroups by conjugation. Thus, for $\pi \in \mathcal{N}$, we have

$$t_i^\pi = t_j^r,$$

where r is an integer coprime to m . Of course, if $m = 2$ then $\mathcal{N}$ will simply act by conjugation as permutations of the n involutory symmetric generators. Now, since by the above elements of $\mathcal{N}$ can be gathered on the left, every element of the progenitor can be represented as πw , where $\pi \in \mathcal{N}$ and w is a word in the symmetric generators. Indeed this representation is unique provided w is simplified so that adjacent symmetric generators are distinct. Thus any additional relator by which we must factor the progenitor to obtain $\mathcal{G}$ must have the form

$$\pi w(t_1, t_2, \dots, t_n),$$

where $\pi \in \mathcal{N}$ and w is a word in $\mathcal{T}$. In the next section we describe how a particular factor group

$$\frac{m^{*n} : \mathcal{N}}{\pi_1 w_1, \dots, \pi_s w_s} \tag{1}$$

may be identified.

3 Manual double coset enumeration

Since in this paper we are only concerned with involutory symmetric generators we restrict our attention to the case $m = 2$. Thus we seek homomorphic images of the progenitor

$$2^{*n} : \mathcal{N},$$

(where $\mathcal{N}$ is now a transitive permutation group on n letters), which act faithfully on $\mathcal{N}$ and on the generators of the free product. It is convenient to identify the n free generators and $\mathcal{N}$ with their respective images. Thus

$$\frac{2^{*n} : \mathcal{N}}{\pi_1 w_1, \dots, \pi_s w_s} \cong \langle \mathcal{N}, \mathcal{T} \mid t_i^2 = 1, t_i^\pi = t_{\pi(i)}, \pi_1 w_1 = \dots = \pi_s w_s = 1 \rangle, \quad (2)$$

where $\pi \in \mathcal{N}$, $t_i \in \mathcal{T}$. Following [10] we are allowing i to stand for the symmetric generator t_i in expressions such as the above relation. By a slight abuse of notation we also allow i to denote the coset $\mathcal{N}t_i$, ij the coset $\mathcal{N}t_i t_j$ etc., and we write, for instance,

$$ij \sim k \text{ to mean } \mathcal{N}t_i t_j = \mathcal{N}t_k.$$

Writing $ij = k$ would be the much stronger statement that $t_i t_j = t_k$. Now since

$$t_i \pi = \pi t_{\pi(i)},$$

(or $i\pi = \pi i^\pi$ as we shall more commonly write), the permutations involved in any element of $\mathcal{G}$ can be gathered on the left. Thus any element of $\mathcal{G}$ can be written as a permutation belonging to $\mathcal{N}$ followed by a word in the symmetric generators. Indeed, as mentioned in the last section, in the case of the progenitor itself this representation is unique provided the obvious cancellations are performed. Thus, if $\mathcal{N}g\mathcal{N}$ is a double coset of $\mathcal{N}$ in $\mathcal{G}$, we have

$$\mathcal{N}g\mathcal{N} = \mathcal{N}\pi w\mathcal{N} = \mathcal{N}w\mathcal{N},$$

where $g = \pi w \in \mathcal{G}$, with $\pi \in \mathcal{N}$, and w is a word in the t_i . We denote this double coset by $[w]$, e.g. $[01]$ denotes the double coset $\mathcal{N}t_0 t_1 \mathcal{N}$. The double coset $\mathcal{N}e\mathcal{N} = \mathcal{N}$, where e is the identity element, is denoted by $[\star]$. Furthermore we define

$$\mathcal{N}^i = \mathcal{C}_{\mathcal{N}}(t_i); \mathcal{N}^{ij} = \mathcal{C}_{\mathcal{N}}(\langle t_i, t_j \rangle) \text{ etc,}$$

single point and two point stabilizers in $\mathcal{N}$ respectively. The coset stabilizing subgroup, $\mathcal{N}^{(w)}$, of $\mathcal{N}$ is given by,

$$\mathcal{N}^{(w)} = \{\pi \in \mathcal{N} : \mathcal{N}w\pi = \mathcal{N}w\},$$

for w a word in the symmetric generators. Clearly $\mathcal{N}^w \leq \mathcal{N}^{(w)}$, and the number of cosets in the double coset $[w] = \mathcal{N}w\mathcal{N}$ is given by $|\mathcal{N}| / |\mathcal{N}^{(w)}|$, since

$$\mathcal{N}w\pi_1 \neq \mathcal{N}w\pi_2 \iff \mathcal{N}w\pi_1\pi_2^{-1} \neq \mathcal{N}w$$

$$\begin{aligned}
&\Longleftrightarrow \pi_1 \pi_2^{-1} \notin \mathcal{N}^{(w)} \\
&\Longleftrightarrow \mathcal{N}^{(w)} \pi_1 \pi_2^{-1} \neq \mathcal{N}^{(w)} \\
&\Longleftrightarrow \mathcal{N}^{(w)} \pi_1 \neq \mathcal{N}^{(w)} \pi_2.
\end{aligned}$$

In order to obtain the index of $\mathcal{N}$ in $\mathcal{G}$ we shall perform a manual double coset enumeration of $\mathcal{G}$ over $\mathcal{N}$; thus we must find all double cosets $[w]$ and work out how many single cosets each of them contains. We shall know that we have completed the double coset enumeration when the set of right cosets obtained is closed under right multiplication. Moreover, the completion test above is best performed by obtaining the orbits of $\mathcal{N}^{(w)}$ on the symmetric generators. We need only identify, for each $[w]$, the double coset to which $\mathcal{N}wt_i$ belongs for one symmetric generator t_i from each orbit. We will decompose the image $\mathcal{G}$ into double cosets $\mathcal{N}g\mathcal{N}$, where $g \in 2^{*n} : \mathcal{N}$ and find a set $\{g_1, g_2, \dots\}$ of elements of $\mathcal{G}$ such that

$$\mathcal{G} = \mathcal{N}g_1\mathcal{N} \cup \mathcal{N}g_2\mathcal{N} \cup \dots$$

But for each i , we have $g_i = \pi_i w_i$, where $\pi_i \in \mathcal{N}$ and w_i is a word in the t_i , and so the double coset decomposition simplifies to

$$\mathcal{G} = \mathcal{N} \cup \mathcal{N}w_2\mathcal{N} \cup \mathcal{N}w_3\mathcal{N} \cup \dots,$$

where w_1 is chosen to be the identity. When the set of relations by which we are factoring is empty this gives the double coset decomposition of the progenitor $2^{*n} : \mathcal{N}$, and in this case there are infinitely many double cosets corresponding to the orbits of $\mathcal{N}$ on the ordered k -tuples of the letters of $\Omega = \{t_1, \dots, t_n\}$ which have no adjacent repetitions, where $k \in \mathbb{N} = \{1, 2, \dots\}$.

We now give two interesting examples to illustrate the process.

Example 1

We consider

$$\mathcal{G} \cong \frac{2^{*6} : L_2(5)}{[(\infty, 0, 1)(2, 4, 3)t_2]^5} \cong \langle \mathcal{N}, \mathcal{T} \mid \mathcal{N} \cong L_2(5), t_i^\pi = t_{\pi(i)}, [(\infty, 0, 1)(2, 4, 3)t_2]^5 = 1 \rangle, \quad (3)$$

Thus $\mathcal{G}$ is the homomorphic image of the split extension $2^{*6} : L_2(5)$ factored by the relator $[(\infty, 0, 1)(2, 4, 3)t_2]^5$ and the action of $\mathcal{N} = L_2(5)$ on the six symmetric generators is given by $x \sim (0, 1, 2, 3, 4)$ and $y \sim (0, \infty)(1, 4)$. Of course at this stage the group could still be infinite, or could collapse to the identity. The given relation

$[\pi t_2]^5 = 1$, where $\pi = (\infty, 0, 1)(2, 4, 3)$, implies

$$\begin{aligned}
&\pi t_2 \pi t_2 \pi t_2 \pi t_2 \pi t_2 = 1 \\
&\Rightarrow \pi^2(\pi)^{-1} t_2 \pi t_2 \pi^3(\pi)^{-2} t_2 \pi^2(\pi)^{-1} t_2 \pi t_2 = 1 \\
&\Rightarrow \pi^2 t_2^\pi t_2 t_2^\pi t_2^\pi t_2 = 1 \\
&\Rightarrow \pi^2 t_4 t_2 t_3 t_4 t_2 = 1 \\
&\Rightarrow t_4 t_2 t_3 t_4 t_2 = \pi.
\end{aligned}$$

Note that, in particular, all conjugates of the relator $\pi^2 t_4 t_2 t_3 t_4 t_2$, under conjugation by $\mathcal{N}$, are also relators. We identify elements of the progenitor $2^{*6} : L_2(5)$

with their images in $\mathcal{G}$ because in the cases that interest us the image of $\mathcal{N}$ will be isomorphic to $\mathcal{N}$ and the images of t_i will be of order 2 and be distinct. The vertices of the Cayley graph of $2^{\star n} : \mathcal{N}$ has the set of right cosets $\{\mathcal{N}w \mid w \text{ a word in the symmetric generators } t_i\}$ as its vertices. Each vertex $\mathcal{N}w$ is joined to the vertex $\mathcal{N}wt_i$, for $i \in \{1, 2, \dots, n\}$. The collapsed Cayley graph is a diagram of the Cayley graph in which each orbit of $\mathcal{N}$ in its action on the vertices by right multiplication is represented by a single node, labelled with the number of vertices that it contains. Lines between these nodes are labelled with integers to indicate how many edges from a vertex of one node lead to vertices of the other. The orbits of $\mathcal{N}$ on the right cosets of $\mathcal{N}$ in $\mathcal{G}$ are, of course, just the double cosets of the form $\mathcal{N}g\mathcal{N}$ and the labels on the nodes indicate how many single cosets a given double coset contains. So in Figure the the node $\mathcal{N}t_\infty t_0 t_1$ is labelled 20, since $\mathcal{N}t_\infty t_0 t_1 = \mathcal{N}t_1 t_2 t_4 = \mathcal{N}t_4 t_3 t_\infty$. Generally, Cayley graphs contain multiple edges and loops. For example, the graph below does not have multiple edges but has two loops, the integer 1+2 at the top of the loop over the node labelled 30 indicates that $[\infty 0 \infty] = [\infty 0]$ and $[\infty 0 1] = [\infty 0]$, with ∞ in the 1-orbit $\{\infty\}$ and 1 in the 2-orbit $\{1, 4\}$. The set of all double cosets $[w] = \mathcal{N}w\mathcal{N}$, coset stabilizing subgroups $\mathcal{N}^{(w)}$, and the number of single cosets each contains are exhibited in Table 1.




Figure 1: The Cayley graph of $L_2(19)$ over $L_2(5)$

Our argument shows that the maximum possible index of $\mathcal{N}$ in $\mathcal{G}$ is $\frac{\mathcal{N}}{\mathcal{N}} + \frac{\mathcal{N}}{\mathcal{N}(\infty)} + \frac{\mathcal{N}}{\mathcal{N}(\infty 0)} + \frac{\mathcal{N}}{\mathcal{N}(\infty 0 1)} = 1 + \frac{\mathcal{N}}{|D_{10}|} + \frac{\mathcal{N}}{|C_2|} + \frac{\mathcal{N}}{|C_3|} = 1 + 6 + 30 + 20 = 57$. Thus $|\mathcal{G}| \leq 57 \times |N| = 57 \times 60 = 3420$. We now show that G is isomorphic to $L_2(19)$. We construct a homomorphism from the progenitor $2^{*6} : L_2(5)$ to $L_2(19)$ by defining $x \equiv (\frac{8\eta+15}{\eta}) = (1, 4, 7, 2, 6)(3, 13, 15, 9, 16)(5, 11, 18, 12, 14)(8, 17, 10, 0, \infty)$, and $y \equiv (\frac{-\eta-1}{2\eta+1}) = (1, 12)(2, 7)(3, 13)(4, 10)(5, 15)(6, 17)(8, 14)(9, 20)(11, 16)(18, 0)$. Since the order of xy is 5, $\mathcal{N} = \langle x, y \rangle \cong A_5$. We now let $t_\infty \equiv (\frac{4\eta+15}{\eta-4}) = (1, 0)(2, 17)(3, 11)(4, \infty)(5, 16)(6, 10)(7, 8)(9, 14)(12, 15)(13, 18)$, and find that $|t_\infty^\mathcal{N}| = 6$ and that $\mathcal{N}$ permutes the six images of t_∞ , by conjugation, as the group $L_2(5)$. Thus

$$\begin{aligned} t_0 &= (1, 5)(2, 14)(3, 19)(4, 17)(6, 7)(8, 20)(9, 10)(11, 15)(12, 18)(13, 16), \\ t_1 &= (1, 2)(3, 15)(4, 11)(5, 6)(7, 10)(8, 17)(9, 18)(12, 14)(13, 20)(16, 19), \\ t_2 &= (1, 11)(2, 19)(3, 20)(4, 6)(5, 14)(7, 18)(8, 15)(9, 13)(10, 17)(12, 16), \\ t_3 &= (1, 7)(2, 12)(3, 14)(4, 18)(5, 11)(6, 20)(8, 13)(9, 17)(10, 19)(15, 16), \\ t_4 &= (1, 8)(2, 4)(3, 9)(5, 13)(6, 14)(7, 12)(10, 16)(11, 18)(15, 17)(19, 20), \\ t_\infty &= (1, 19)(2, 17)(3, 11)(4, 20)(5, 16)(6, 10)(7, 8)(9, 14)(12, 15)(13, 18), \end{aligned}$$

where, under conjugation, we have:

$$x: (t_1, t_2, t_3, t_4, t_0)(t_\infty) \text{ and}$$

$$y: (t_\infty, t_0)(t_1, t_4)(t_2)(t_3).$$

Since $\mathcal{N} \cong A_5$ is maximal in $L_2(19)$ and $t_\infty \notin \mathcal{N}$, we can conclude that $L_2(19)$ is a homomorphic image of the progenitor $2^{*6} : L_2(5)$; it remains to check that the additional relation $[\pi t_2]^5 = 1 \iff t_4 t_2 t_3 t_4 t_2 = \pi = (\infty, 0, 1)(2, 4, 3) = ((x * y)^{(x^2 * y * x^3 * y * x^3)}) \iff ((x * y)^{(x^2 * y * x^3 * y * x^3)} * t_2)^5 = 1$ holds. We readily calculate that

$$\begin{aligned} t_4 t_2 t_3 t_4 t_2 &= (1, 17, 14)(3, 11, 15)(4, 12, 19)(5, 8, 9)(6, 7, 10)(16, 20, 18) \\ &= ((x * y)^{(x^2 * y * x^3 * y * x^3)}), \end{aligned}$$

which acts as $(\infty, 0, 1)(2, 4, 3)$, by conjugation, on the six symmetric generators. This shows that $L_2(19)$ is an image of G ; but $|G| \leq 57 \times 60 = |L_2(19)|$, and so the equality holds and $G \cong L_2(19)$.

The Cayley graph of $L_2(19)$ over $L_2(5)$ suggests that every element of $L_2(19)$ can be written as a permutation, of $L_2(5)$ on six letters followed by a word, in term of the symmetric generators, of length at most three.

We now give an example of a progenitor where the homomorphic image $\mathcal{G}$ does not possess a symmetric generating set. In fact, the symmetric generators will generate a subgroup of $\mathcal{G}$ of index 2.

Example 2

We consider

$$\mathcal{G} \cong \frac{2^{*3} : S_3}{[(0, 1, 2)t_0]^{10}, [(0, 1)t_0]^6} \cong \langle \mathcal{N}, \mathcal{T} \mid \mathcal{N} \cong S_3, t_i^\pi = t_{\pi(i)}, [(0, 1, 2)t_0]^{10} = 1, [(0, 1)t_0]^6 \rangle, \quad (4)$$

Label $[w]$	Coset stabilizing subgroup $\mathcal{N}^{(w)}$	Number of cosets
$[\star]$	Since $\mathcal{N}$ is transitive on T	1
$[\infty]$	$\mathcal{N}^{(\infty)} = N^\infty = \langle x, (1, 3)(0, 4) \rangle \cong D_{10}$, has orbits $\{\infty\}$ and $\{0, 1, 2, 3, 4\}$ on T .	6
$[\infty 0]$	$\mathcal{N}^{\infty 0} = N^{(\infty 0)} = \langle (1, 4)(2, 3) \rangle \cong C_2$, has orbits $\{0\}$, $\{\infty\}$, $\{1, 4\}$ and $\{2, 3\}$.	30
$[\infty 00] = [\infty]$	We show that $\infty 0 \infty \sim \infty 0$. Firstly, $\infty.01401 = \infty.(1, 4, 0)(2, 3, \infty)$ $\implies \infty.01401 = (1, 4, 0)(2, 3, \infty)2$ $\implies \infty.014 = (1, 4, 0)(2, 3, \infty)210$ Thus $\infty 014 \sim 210$ and $210 = (1, 2, 0)(3, \infty, 4)12 \implies \infty 014 \sim 12$. So $\infty 01 \sim 124$. Secondly, $\infty.04104 = \infty.(1, 4, 0)(2, \infty, 3)$ $\implies \infty.04104 = (1, 4, 0)(2, \infty, 3)3$. Thus $\infty 041 \sim 340 \sim 43$, since $34034 = (1, 2, \infty)(3, 0, 4)$ $\implies \infty 04 \sim 431$. Now $\infty 0 \infty = \infty 01.1 \infty \sim 124.1 \infty =$ $12.41 \infty 41.14 = 12.(1, 4, \infty)(2, 3, 0).14 \sim 4314 \sim \infty 044 \sim \infty 0$. and $\infty 02 \infty 0 = (1, 3, 4)(2, 0, \infty) \implies \infty 02 \sim 0 \infty$ $\infty 01 = \infty 03.31 \sim 0 \infty 31 \sim 0(1, 3, \infty)(2, 0, 4)3 \infty)3 \infty \sim 43 \infty$. Thus $\infty 01 \sim 43 \infty \sim 124$	
$[\infty 0 \infty] = [\infty 0]$		
$[\infty 02] = [\infty 0]$		
$[\infty 01]$	$\mathcal{N}^{(\infty 01)} = \langle (1, 4, \infty)(2, 3, 0) \rangle \cong C_3$, has orbits $\{1, 4, \infty\}$ and $\{2, 3, 0\}$ on $\mathcal{T}$. Now $\infty 012 \sim \infty.(1, 0, 2)(3, 4, \infty)10 \sim 310$.	20
$[\infty 012] = [\infty 01]$		
$[\infty 011] = [\infty 0]$		

Table 1: The double cosets $[w] = \mathcal{N}w\mathcal{N}$ in $L_2(19)$.

The group $\mathcal{G}$, given above, is isomorphic to $5^2 : D_6$.

Manual double coset enumeration of $5^2 : D_6$ over S_3 :

We first note that the double coset $[0] = \mathcal{N}t_0\mathcal{N} = \{\mathcal{N}t_0, \mathcal{N}t_1, \mathcal{N}t_2\}$ and the double coset $[01] = \mathcal{N}t_0t_1\mathcal{N} = \{\mathcal{N}t_0t_1, \mathcal{N}t_0t_2, \mathcal{N}t_1t_0, \mathcal{N}t_1t_2, \mathcal{N}t_2t_0, \mathcal{N}t_2t_1\}$. In order to compute the remaining double cosets, we first examine our relations. The relation $[\pi t_0]^{10} = 1$, where $\pi = (0, 1, 2)$, gives

$$\begin{aligned} & \pi t_0 \pi t_0 \pi t_0 \pi t_0 \pi t_0 \pi t_0 \pi t_0 \pi t_0 \pi t_0 \pi t_0 = 1 \\ \implies & \pi \pi \pi^{-1} t_0 \pi t_0 \pi \pi \pi^{-1} t_0 \pi t_0 \pi \pi \pi^{-1} t_0 \pi t_0 \pi \pi \pi^{-1} t_0 \pi t_0 \pi \pi \pi^{-1} t_0 \pi t_0 = 1 \\ \implies & \pi^{-1} t_1 t_0 \pi^{-1} t_1 t_0 \pi^{-1} t_1 t_0 \pi^{-1} t_1 t_0 \pi^{-1} t_1 t_0 \pi^{-1} t_1 t_0 = 1 \\ \implies & \pi^{-1} \pi^{-1} \pi t_1 t_0 \pi^{-1} t_1 t_0 \pi^{-1} \pi^{-1} \pi t_1 t_0 \pi^{-1} t_1 t_0 \pi^{-1} \pi^{-1} \pi t_1 t_0 = 1 \\ \implies & \end{aligned}$$

$$\pi t_0 t_2 t_1 t_0 t_2 t_1 t_0 t_2 t_1 t_0 = 1$$

(Note that $[\pi t_0]^{10} = 1$ can be written as

$$\pi^{10} t_0^9 t_0^8 t_0^7 t_0^6 t_0^5 t_0^4 t_0^3 t_0^2 t_0 = 1).$$

Thus $(0, 1, 2)t_0 t_2 t_1 t_0 t_2 t_1 t_0 t_2 t_1 t_0 = 1$ and

$$(0, 1, 2)t_0 t_2 t_1 t_0 t_2 = t_0 t_1 t_2 t_0 t_1. \quad (5)$$

Also, the relation $[(0, 1)t_0]^6 = 1 \implies (0, 1)t_0(0, 1)t_0(0, 1)t_0(0, 1)t_0(0, 1)t_0(0, 1)t_0 = t_0^{(0,1)} t_0^{(0,1)} t_0^{(0,1)} t_0^{(0,1)} t_0 = t_1 t_0 t_1 t_0 t_1 t_0 = 1 \implies$

$$t_0 t_1 t_0 = t_1 t_0 t_1 \quad (6)$$

Now $t_0 t_1 t_2 t_0 t_2 t_1 t_0 = t_0 t_1 . t_2 t_0 t_2 . t_1 t_0 = t_0 t_1 . t_0 t_2 t_0 . t_1 t_0 = t_0 t_1 t_0 . t_2 . t_0 t_1 t_0 = t_1 t_0 t_1 . t_2 . t_1 t_0 t_1 = t_1 t_0 . t_1 t_2 t_1 . t_0 t_1 = t_1 t_0 . t_2 t_1 t_2 . t_0 t_1$. This gives

$$t_0 t_1 t_2 t_0 t_2 t_1 t_0 = t_1 t_0 t_2 t_1 t_2 t_0 t_1 \quad (7)$$

Now $\mathcal{N}^{01} = N^{(01)} \geq e$, has orbits $\{0\}$, $\{1\}$ and $\{2\}$. Thus we must consider the double cosets $[010]$, $[011]$ and $[012]$. However, $[011] = [0]$ and $[012] = \mathcal{N}t_0 t_1 t_2 \mathcal{N} = \{\mathcal{N}t_0 t_1 t_2, \mathcal{N}t_0 t_2 t_1, \mathcal{N}t_1 t_0 t_2, \mathcal{N}t_1 t_2 t_0, \mathcal{N}t_2 t_0 t_1, \mathcal{N}t_2 t_1 t_0\}$. Moreover, (by) $010 = 101 \implies |\mathcal{N}^{(010)}| \geq |< (0, 1) >| = 2$. Thus the double coset $[010]$ contains $\frac{|\mathcal{N}|}{|\mathcal{N}^{(010)}|} = 3$ single cosets. Now $N^{(010)} \geq e < (0, 1) > \cong C_2$ has orbits $\{0, 1\}$ and $\{2\}$ on $\mathcal{T}$ and $0102 \sim 1012$. Thus the double coset $[0102]$ contains three single cosets. $\mathcal{N}^{(0102)} = e < (0, 1) > \cong C_2$ has orbits $\{0, 1\}$ and $\{2\}$ on $\mathcal{T}$ and $020 = 202 \implies 01020 = 01202$. Thus $[01020] = [01202]$ and the double coset has six single cosets. At this point we have the double cosets $[012]$ and $[01202]$ left to be considered.

Now $N^{(012)} = e < e >$ has orbits $\{0\}$, $\{1\}$ and $\{2\}$ on $\mathcal{T}$. But $121 = 212 \implies 0121 = 0212 \implies$ the coset 0121 has two names. Thus $[0121]$ contains 3 single cosets. Now $N^{(0121)} \geq e < (1, 2) > \cong C_2$ has orbits $\{0\}$ and $\{1, 2\}$ on $\mathcal{T}$ and $01210 = 02120 \implies$ the coset 01210 has two names. Thus $[01210]$ contains three single cosets and $N^{(01210)} \geq e < (1, 2) > \cong C_2$ has orbits $\{0\}$ and $\{1, 2\}$ on $\mathcal{T}$ with $012101 = 012010 \implies [012101] = [012010]$ and since

$(0, 1, 2)t_0t_2t_1t_0t_2 = t_0t_1t_2t_0t_1 \implies t_0t_1t_2t_0t_1t_0 = (0, 1, 2)t_0t_2t_1t_0t_2t_0$, and therefore $012010 \sim 021020$, the double coset $[012010]$ contains three single cosets. We now have the double cosets $[0120]$ and $[01202]$ that must be considered. Now $N^{(0120)} \geq \langle e \rangle$ has orbits $\{0\}$, $\{1\}$ and $\{2\}$ on $\mathcal{T}$. Because of the relation $01201 = (0, 1, 2)02102$, the coset 01201 has two names. Thus $[01201]$ contains three single cosets. Since $N^{(01201)} \geq \langle (1, 2) \rangle \cong C_2$ has orbits $\{0\}$, $\{1, 2\}$ on $\mathcal{T}$ and the double coset $[012010]$ has already been investigated. We are now left with the double coset $[01202]$. From above, $[01202]$ contains 6 single cosets. Since $N^{(01202)} \geq \langle e \rangle$ has orbits $\{0\}$, $\{1\}$ and $\{2\}$ on $\mathcal{T}$ and $01201 = (0, 1, 2)02102 \implies 0120 = (0, 1, 2)021021 \implies 012021 = (0, 1, 2)02102121 \implies 012021 = (0, 1, 2)021012$, the coset 012021 has two names. Thus the double coset $[012021]$ contains 3 single cosets. $N^{(012021)} \geq \langle (1, 2) \rangle \cong C_2$ has orbits $\{0\}$ and $\{1, 2\}$ on $\mathcal{T}$. From above, $012021 = (0, 1, 2)021012$, $0120210 = (0, 1, 2)0210120 = (0, 1, 2)0201020 = (0, 1, 2)2021020 = (0, 1, 2)2021202 = (0, 1, 2)2012102$. Thus $0120210 = (0, 1, 2)0210120 = (0, 1, 2)2012102$. Then, conjugation by $(0, 1)$ gives, $1021201 = (1, 0, 2)1201021 = (1, 0, 2)2102012$ and $0120210 = 1021201$ (page). Hence $0120210 = (0, 1, 2)0210120 = (0, 1, 2)2012102 = 1021201 = (0, 1, 2)1201021 = (0, 1, 2)2102012$. $[0120210]$ contains 1 coset and $N^{(0120210)} \geq \langle (1, 2), (0, 1, 2) \rangle \cong S_3$ is transitive on $\mathcal{T}$.




Figure 2: The Cayley graph of $5^2 : D_6$ over S_3




Figure 3: 7-point Projective Plane.

We now present our main problem. A presentation for the progenitor $2^{*(7+7)} : PGL_2(7)$, is given by :

$$\begin{aligned} &< x, y, t | x^7, y^2, t^2, (x^{-1} * t)^2, (y * x)^3, t * x^{-1} * y \\ &* x * t * y, x^2 * y * x^3 * y * x^{-4} * y * x^{-4} * y * x, s^2, (s^3, y), (s^4, x * y) >, \\ &\text{where } N = PGL_2(7) \cong < x, y, t | x^7, y^2, t^2, (x^{-1} * t)^2, (y * x)^3, \\ &t * x^{-1} * y * x * t * y, x^2 * y * x^3 * y * x^{-4} * y * x^{-4} * y * x >; \end{aligned}$$

and the action of N on the symmetric generators is given by

$$\begin{aligned} x &\sim (0, 1, 2, 3, 4, 5, 6)(0, 6, 5, 4, 3, 2, 1), \\ y &\sim (2, 6)(4, 5)(0, 3)(5, 6), \\ t &\sim (0, 0)(1, 1)(2, 2)(3, 3)(4, 4)(5, 5)(6, 6). \end{aligned}$$

We factor the progenitor by the following relations

$$t = \mathbf{000} \text{ and } y = \mathbf{1111}$$

to get the finite homomorphic image G . Thus

$$G \cong \frac{2^{*(7+7)} : PGL_2(7)}{t = \mathbf{000}, y = (\mathbf{01})^2}. \quad (8)$$

Label $[w]$	Coset stabilizing subgroup $\mathcal{N}^{(w)}$	Number of cosets
$[\star]$	Since $\mathcal{N}$ is transitive on T	1
$[0]$	$\mathcal{N}^{(0)} = N^0 \geq \langle (1, 2) \rangle \cong C_2$, has orbits $\{0\}$ and $\{1, 2\}$ on $\mathcal{T}$.	6
$[01]$	$\mathcal{N}^{01} = N^{(01)} \geq \langle e \rangle$, has orbits $\{0\}$, $\{1\}$ and $\{2\}$. Moreover, $010 = 101 \implies N^{(010)} \geq \langle (0, 1) \rangle = 2$	6
$[010]$	$N^{(010)} \geq \langle (0, 1) \rangle \cong C_2$ has orbits $\{0, 1\}$ and $\{2\}$ on $\mathcal{T}$ and $0102 \sim 1012$. Thus the coset 0102 has two names. This implies	3
$[0102]$	$N^{(0102)} = \langle (0, 1) \rangle \cong C_2$ has orbits $\{0, 1\}$ and $\{2\}$ on $\mathcal{T}$ and $020 = 202 \implies 01020 = 01202$.	3
$[01021] = [01202]$		6
$[012]$	$N^{(012)} = \langle e \rangle$ has orbits $\{0\}$, $\{1\}$ and $\{2\}$ on $\mathcal{T}$.	6
$[0120]$	Now $121 = 212 \implies 0121 = 0212 \implies$ the coset 0121 has two names,	6
$[0121]$	$N^{(0121)} \geq \langle (1, 2) \rangle \cong C_2$ has orbits $\{0\}$ and $\{1, 2\}$ on $\mathcal{T}$ and $01210 = 02120 \implies$ the coset 01210 has two names.	3
$[01210]$	$N^{(01210)} \geq \langle (1, 2) \rangle \cong C_2$ has orbits $\{0\}$ and $\{1, 2\}$ on $\mathcal{T}$ and $012101 = 012010$ and the relation $01201 = (0, 1, 2)02102$ implies $012010 = (0, 1, 2)021020$ and so the coset 012010 has two names.	3
$[012101] = [012010]$	$N^{(0120)} \geq \langle e \rangle$ has orbits $\{0\}$, $\{1\}$ and $\{2\}$ on $\mathcal{T}$. Because of the relation $01201 = (0, 1, 2)02102$, the coset 01201 has two names.	3
$[01201]$	$N^{(01201)} \geq \langle (1, 2) \rangle \cong C_2$ has orbits $\{0\}$, $\{1, 2\}$ on $\mathcal{T}$ and the relation $01201 = (0, 1, 2)02102$ implies $012010 = (0, 1, 2)021020$ and so the coset 012010 has two names.	3
$[012010]$	$N^{(01202)} \geq \langle e \rangle$ has orbits $\{0\}$, $\{1\}$ and $\{2\}$ on $\mathcal{T}$ and the relation $01201 = (0, 1, 2)02102 \implies 0120 = (0, 1, 2)021021 \implies 012021 = (0, 1, 2)02102121 \implies 012021 = (0, 1, 2)021012$ implies that the coset 012021 has two names.	3
$[012021]$	$N^{(012021)} \geq \langle (1, 2) \rangle \cong C_2$ has orbits $\{0\}$ and $\{1, 2\}$ on $\mathcal{T}$. From above, $012021 = (0, 1, 2)021012$, $0120210 = (0, 1, 2)0210120 = (0, 1, 2)0201020 = (0, 1, 2)0201020 = (0, 1, 2)0201202 = (0, 1, 2)0201202$. Thus $0120210 = (0, 1, 2)0210120 = (0, 1, 2)0201202$. Then, conjugation by $(0, 1)$ gives, $1021201 = (1, 0, 2)1201021 = (1, 0, 2)2102012$ and $0120210 = 1021201$ (page). Hence $0120210 = (0, 1, 2)0210120 = (0, 1, 2)0201202 = 1021201 = (0, 1, 2)1201021 = (0, 1, 2)2102012$.	3
$[0120210]$	$N^{(0120210)} \geq \langle (1, 2), (0, 1, 2) \rangle \cong S_3$ is transitive on $\mathcal{T}$.	1

Table 2: The double cosets $[w] = \mathcal{N}w\mathcal{N}$ in $5^2 : D_6$.

The index of $PGL_2(7)$ in G is 36. It turns out that $G \cong U_3(3) : 2$ (see [9]).

4 Manual double coset enumeration of $U_3(3) : 2$ over $PGL_2(7)$

$$t = \mathbf{000} \implies \mathbf{00} = t\mathbf{0} \implies \mathbf{00} \sim \mathbf{0}.$$

$$y = (\mathbf{01})^2 = \mathbf{0101} \implies \mathbf{01} \sim \mathbf{10}. \text{ Define } \pi_{\mathbf{000}} = t = (\mathbf{0}, 0)(\mathbf{1}, 1)(\mathbf{2}, 2)(\mathbf{3}, 3)(\mathbf{4}, 4)(\mathbf{5}, 5)(\mathbf{6}, 6),$$

$$\text{and } \sigma_{\mathbf{01}} = \mathbf{0101} = y = (\mathbf{2}, \mathbf{6})(\mathbf{4}, \mathbf{5})(\mathbf{0}, 3)(\mathbf{5}, 6).$$

We note that

$$\pi_{\mathbf{050}} = (\mathbf{1}, 6)(\mathbf{2}, 3)(\mathbf{3}, 1)(\mathbf{4}, 4)(\mathbf{5}, 0)(\mathbf{6}, 2)(\mathbf{0}, 5) \text{ and}$$

$$\pi_{\mathbf{515}} = (\mathbf{1}, 5)(\mathbf{2}, 4)(\mathbf{3}, 1)(\mathbf{4}, 0)(\mathbf{5}, 2)(\mathbf{6}, 3)(\mathbf{0}, 6). \text{ Then}$$

$$\mathbf{01} = \mathbf{0551}$$

$$= \mathbf{05.515.5}$$

$$= \mathbf{050.0.515.5}$$

$$= (\mathbf{1}, 6)(\mathbf{2}, 3)(\mathbf{3}, 1)(\mathbf{4}, 4)(\mathbf{5}, 0)(\mathbf{6}, 2)(\mathbf{0}, 5). \mathbf{0.1.5}(\mathbf{2}, 4)(\mathbf{3}, 1)(\mathbf{4}, 0)(\mathbf{5}, 2)(\mathbf{6}, 3)(\mathbf{0}, 6). \mathbf{5}$$

$$= \mathbf{05.515.5}$$

$$= \mathbf{050.0.515.5}$$

$$= (\mathbf{1}, 6)(\mathbf{2}, 3)(\mathbf{3}, 1)(\mathbf{4}, 4)(\mathbf{5}, 0)(\mathbf{6}, 2)(\mathbf{0}, 5)(\mathbf{1}, 5)(\mathbf{2}, 4)(\mathbf{3}, 1)(\mathbf{4}, 0)(\mathbf{5}, 2)(\mathbf{6}, 3)(\mathbf{0}, 6). \mathbf{65.}$$

$$\text{Thus } \mathbf{01} = (\mathbf{0}, \mathbf{1})(\mathbf{2}, \mathbf{6}), (\mathbf{5}, \mathbf{4})(\mathbf{0}, 2, 3, 4)(\mathbf{5}, 6). \mathbf{65.}$$

$$\text{So } [\mathbf{01}]^2 = \mathbf{0101}$$

$$= (\mathbf{0}, \mathbf{1})(\mathbf{2}, \mathbf{6}, \mathbf{5}, \mathbf{4})(\mathbf{0}, 2, 3, 4)(\mathbf{5}, 6). \mathbf{65.}(\mathbf{0}, \mathbf{1})(\mathbf{2}, \mathbf{6}, \mathbf{5}, \mathbf{4})(\mathbf{0}, 2, 3, 4)(\mathbf{5}, 6). \mathbf{65}$$

$$= (\mathbf{2}, \mathbf{5})(\mathbf{4}, \mathbf{6})(\mathbf{0}, 3)(\mathbf{2}, 4).$$

$$\implies \mathbf{10} = (\mathbf{2}, \mathbf{5})(\mathbf{4}, \mathbf{6})(\mathbf{0}, 3)(\mathbf{2}, 4)\mathbf{01}.$$

$$\text{Define } \tau_{\mathbf{01}} = \mathbf{0101} = (\mathbf{2}, \mathbf{5})(\mathbf{4}, \mathbf{6})(\mathbf{0}, 3)(\mathbf{2}, 4).$$

Similarly,

$$\mathbf{10} = \mathbf{1660}$$

$$= \mathbf{16.606.6}$$

$$= \mathbf{161.1.606.6}$$

$$= (\mathbf{1}, 6)(\mathbf{2}, 0)(\mathbf{3}, 1)(\mathbf{4}, 2)(\mathbf{5}, 3)(\mathbf{6}, 4)(\mathbf{0}, 5). \mathbf{1.1.5}(\mathbf{2}, 2)(\mathbf{3}, 1)(\mathbf{4}, 3)(\mathbf{5}, 4)(\mathbf{6}, 0)(\mathbf{0}, 6). \mathbf{6}$$

$$= (\mathbf{1}, 6)(\mathbf{2}, 0)(\mathbf{3}, 1)(\mathbf{4}, 2)(\mathbf{5}, 3)(\mathbf{6}, 4)(\mathbf{0}, 5)(\mathbf{1}, 5)(\mathbf{2}, 2)(\mathbf{3}, 1)(\mathbf{4}, 3)(\mathbf{5}, 4)(\mathbf{6}, 0)(\mathbf{0}, 6). \mathbf{56}$$

$$= (\mathbf{0}, \mathbf{1})(\mathbf{2}, \mathbf{6}, \mathbf{5}, \mathbf{4})(\mathbf{0}, 2, 3, 4)(\mathbf{5}, 6). \mathbf{56.}$$

$$\text{Now } \mathbf{1} \sim \mathbf{1000} \sim \mathbf{0100} \sim \mathbf{0.1010.010} \sim \mathbf{0(3, 0)(5, 6)(2, 6)(4, 5)010} \sim \mathbf{3010} \sim$$

$$\mathbf{30(2, 5)(4, 6)(0, 3)(2, 4)01} \sim \mathbf{3301} \sim \mathbf{301.}$$

$$\implies \mathbf{11} \sim \mathbf{30}. \text{ Hence } \mathbf{11} \sim \mathbf{30} \sim \mathbf{03} \sim \mathbf{03} \sim \mathbf{30}.$$

$$\pi_{\mathbf{000}}\mathbf{1} = \mathbf{1}\pi_{\mathbf{000}}$$

$$= \mathbf{1000} = \tau_{\mathbf{01}}\mathbf{0100}$$

$$= \tau_{\mathbf{01}}\mathbf{0}\sigma_{\mathbf{10}}\mathbf{010}$$

$$= \tau_{\mathbf{01}}\sigma_{\mathbf{10}}\mathbf{3010}$$

$$= \tau_{\mathbf{01}}\sigma_{\mathbf{10}}\mathbf{30}\tau_{\mathbf{01}}\mathbf{01}$$

$$= \tau_{\mathbf{01}}\sigma_{\mathbf{10}}\pi_{\mathbf{303}}\mathbf{310} = \tau_{\mathbf{01}}\sigma_{\mathbf{10}}\mathbf{30}\tau_{\mathbf{01}}\mathbf{01}$$

$$= \tau_{\mathbf{01}}\sigma_{\mathbf{10}}\pi_{\mathbf{303}}\mathbf{3}\tau_{\mathbf{01}}\mathbf{01}$$

$$= \tau_{\mathbf{01}}\sigma_{\mathbf{10}}\mathbf{30}\tau_{\mathbf{01}}\mathbf{01} = \tau_{\mathbf{01}}\sigma_{\mathbf{10}}\pi_{\mathbf{303}}\tau_{\mathbf{01}}\mathbf{301}$$

$$= (\mathbf{2}, \mathbf{5})(\mathbf{4}, \mathbf{6})(\mathbf{0}, 3)(\mathbf{2}, 4).(\mathbf{0}, \mathbf{3})(\mathbf{5}, \mathbf{6})(\mathbf{6}, 2)(\mathbf{4}, 5).$$

$(\mathbf{0}, 3)(\mathbf{1}, 1)(\mathbf{2}, 6)(\mathbf{3}, 0)(\mathbf{4}, 5)(\mathbf{5}, 2)(\mathbf{6}, 4).(\mathbf{2}, \mathbf{5})(\mathbf{4}, \mathbf{6})(0, 3)(2, 4)\mathbf{301}.$
 $= (\mathbf{0}, 3, \mathbf{3}, 0)(\mathbf{1}, 1)(\mathbf{2}, 2, \mathbf{6}, 5, \mathbf{4}, \mathbf{4}, \mathbf{5}, 6)\mathbf{301}.$
 Thus $1 = (\mathbf{2}, \mathbf{6})(\mathbf{4}, \mathbf{5})(0, 3)(5, 6)\mathbf{301}.$

5 Manual double coset enumeration of $U_3(3) : 2$ over $PGL_2(7)$

We first note that $\mathcal{N}t_0\mathcal{N}$ contains fourteen single cosets, namely, $\mathcal{N}t_0, \mathcal{N}t_1, \dots, \mathcal{N}t_6, \mathcal{N}t_0, \mathcal{N}t_1, \dots, \mathcal{N}t_6$. In order to compute the remaining double coset $\mathcal{N}w\mathcal{N}$, where w is a word in the t_i 's, we first analyze our relations.

The given relation $t = \mathbf{000}$ gives $\mathbf{00} = t\mathbf{0}$. Thus $\mathbf{00} \sim \mathbf{0}$.

Now $y = (\mathbf{01})^2 = \mathbf{0101} \implies \mathbf{01} \sim \mathbf{10}$.

Define $\pi_{\mathbf{00}} = t = (\mathbf{0}, 0)(\mathbf{1}, 1)(\mathbf{2}, 2)(\mathbf{3}, 3)(\mathbf{4}, 4)(\mathbf{5}, 5)(\mathbf{6}, 6)$,
 and $\mathbf{0101} = y = (\mathbf{2}, \mathbf{6})(\mathbf{4}, \mathbf{5})(0, 3)(5, 6) \implies [\mathbf{0101}]^a = y^a$. Thus

$$\mathbf{1010} = (\mathbf{3}, \mathbf{0})(\mathbf{5}, \mathbf{6})(2, 6)(4, 5). \quad (9)$$

Now

$$\begin{aligned}
 \mathbf{01} &= \mathbf{0551} \\
 &= \mathbf{05.515.5} \\
 &= \mathbf{050.0.515.5} \\
 &= \pi_{\mathbf{05}}.\mathbf{0}.\pi_{\mathbf{15}}.\mathbf{5} \\
 &= \pi_{\mathbf{05}}.\pi_{\mathbf{15}}.\mathbf{0}^{\pi_{\mathbf{15}}}.\mathbf{5} \\
 &= (\mathbf{1}, 6)(\mathbf{2}, 3)(\mathbf{3}, 1)(\mathbf{4}, 4)(\mathbf{5}, 0)(\mathbf{6}, 2)(\mathbf{0}, 5).(\mathbf{1}, 5)(\mathbf{2}, 4)(\mathbf{3}, 1)(\mathbf{4}, 0)(\mathbf{5}, 2)(\mathbf{6}, 3)(\mathbf{0}, 6).65 \\
 \text{Thus } \mathbf{01} &= (\mathbf{0}, \mathbf{1})(\mathbf{2}, \mathbf{6}, \mathbf{5}, \mathbf{4})(0, 2, 3, 4)(5, 6).65.
 \end{aligned}$$

Similarly,

$$\begin{aligned}
 \mathbf{10} &= \mathbf{1660} \\
 &= \mathbf{16.606.6} \\
 &= \mathbf{161.1.606.6} \\
 &= \pi_{\mathbf{16}}.\mathbf{1}.\pi_{\mathbf{06}}.\mathbf{6} \\
 &= \pi_{\mathbf{16}}.\pi_{\mathbf{06}}.\mathbf{1}^{\pi_{\mathbf{06}}}.\mathbf{6} \\
 &= (\mathbf{1}, 6)(\mathbf{2}, 0)(\mathbf{3}, 1)(\mathbf{4}, 2)(\mathbf{5}, 3)(\mathbf{6}, 4)(\mathbf{0}, 5)(\mathbf{1}, 5)(\mathbf{2}, 2)(\mathbf{3}, 1)(\mathbf{4}, 3)(\mathbf{5}, 4)(\mathbf{6}, 0)(\mathbf{0}, 6).56 \\
 &= (\mathbf{0}, \mathbf{1})(\mathbf{2}, \mathbf{6}, \mathbf{5}, \mathbf{4})(0, 2, 3, 4)(5, 6).56.
 \end{aligned}$$

Thus $\mathcal{N}t_0t_1 = \mathcal{N}t_6t_5 = \mathcal{N}t_65t_6$.

Moreover, $[\mathbf{01}]^2 = \mathbf{0101}$

$$\begin{aligned}
 &= (\mathbf{0}, \mathbf{1})(\mathbf{2}, \mathbf{6}, \mathbf{5}, \mathbf{4})(0, 2, 3, 4)(5, 6).65.(\mathbf{0}, \mathbf{1})(\mathbf{2}, \mathbf{6}, \mathbf{5}, \mathbf{4})(0, 2, 3, 4)(5, 6).65 \\
 &= (\mathbf{0}, \mathbf{1})(\mathbf{2}, \mathbf{6}, \mathbf{5}, \mathbf{4})(0, 2, 3, 4)(5, 6).(\mathbf{0}, \mathbf{1})(\mathbf{2}, \mathbf{6}, \mathbf{5}, \mathbf{4})(0, 2, 3, 4)(5, 6).[65]^{(\mathbf{0}, \mathbf{1})(\mathbf{2}, \mathbf{6}, \mathbf{5}, \mathbf{4})(0, 2, 3, 4)(5, 6)}.65 \\
 &= (\mathbf{2}, \mathbf{5})(\mathbf{4}, \mathbf{6})(0, 3)(2, 4)56.65 \\
 &= (\mathbf{2}, \mathbf{5})(\mathbf{4}, \mathbf{6})(0, 3)(2, 4). \text{ Thus}
 \end{aligned}$$

$$[\mathbf{01}]^2 = (\mathbf{2}, \mathbf{5})(\mathbf{4}, \mathbf{6})(0, 3)(2, 4). \quad (10)$$

Also,

$$\mathbf{10} = (\mathbf{2}, \mathbf{5})(\mathbf{4}, \mathbf{6})(0, 3)(2, 4)\mathbf{01}. \quad (11)$$

Hence

$$\mathbf{01} \sim \mathbf{10} \sim 56 \sim 65. \quad (12)$$

From (8) and (9) above, define $\sigma_{01} = \mathbf{0101} = y = (\mathbf{2}, \mathbf{6})(\mathbf{4}, \mathbf{5})(0, 3)(5, 6)$ and $\tau_{01} = \mathbf{0101} = (\mathbf{2}, \mathbf{5})(\mathbf{4}, \mathbf{6})(0, 3)(2, 4)$.

Now

$$\begin{aligned} \pi_{00}1 &= \mathbf{1}\pi_{00} \\ &= \mathbf{1000} = \mathbf{10.00} = \tau_{01}\mathbf{01.00} \\ &= \tau_{01}\mathbf{0.10.0} \\ &= \tau_{01}\mathbf{0}\sigma_{10}\mathbf{010} \\ &= \tau_{01}\mathbf{0}\sigma_{10}0^{\sigma_{10}}\mathbf{10} \\ &= \tau_{01}\sigma_{10}\mathbf{0}^{\sigma_{10}}\mathbf{010} \\ &= \tau_{01}\sigma_{10}\mathbf{3010} \\ &= \tau_{01}\sigma_{10}\pi_{30}\mathbf{310} \\ &= \tau_{01}\sigma_{10}\pi_{30}\mathbf{3}\tau_{01}\mathbf{01} \\ &= \tau_{01}\sigma_{10}\pi_{30}\tau_{01}\mathbf{3}^{\tau_{01}}\mathbf{01} \\ &= \tau_{01}\sigma_{10}\pi_{30}\tau_{01}\mathbf{301} \end{aligned}$$

Hence $\pi_{00}1 = \tau_{01}\sigma_{10}\pi_{30}\tau_{01}\mathbf{301}$.

$$\begin{aligned} \text{So } 1 &= \pi_{00}^{-1}\tau_{01}\sigma_{10}\pi_{30}\tau_{01}\mathbf{301} \\ &= (\mathbf{1}, \mathbf{1})(\mathbf{2}, \mathbf{2})(\mathbf{3}, \mathbf{3})(\mathbf{4}, \mathbf{4})(\mathbf{5}, \mathbf{5})(\mathbf{6}, \mathbf{6})(\mathbf{7}, \mathbf{7}).(\mathbf{2}, \mathbf{5})(\mathbf{4}, \mathbf{6})(0, 3)(2, 4).(\mathbf{0}, \mathbf{3})(\mathbf{5}, \mathbf{6})(6, 2)(4, 5). \\ &\quad (\mathbf{0}, \mathbf{3})(\mathbf{1}, \mathbf{1})(\mathbf{2}, \mathbf{6})(\mathbf{3}, \mathbf{0})(\mathbf{4}, \mathbf{5})(\mathbf{5}, \mathbf{2})(\mathbf{6}, \mathbf{4}).(\mathbf{2}, \mathbf{5})(\mathbf{4}, \mathbf{6})(0, 3)(2, 4)\mathbf{301}. \\ &= (\mathbf{1}, \mathbf{1})(\mathbf{2}, \mathbf{2})(\mathbf{3}, \mathbf{3})(\mathbf{4}, \mathbf{4})(\mathbf{5}, \mathbf{5})(\mathbf{6}, \mathbf{6})(\mathbf{7}, \mathbf{7}).(\mathbf{0}, \mathbf{3}, \mathbf{3}, \mathbf{0})(\mathbf{1}, \mathbf{1})(\mathbf{2}, \mathbf{2}, \mathbf{6}, \mathbf{5}, \mathbf{4}, \mathbf{4}, \mathbf{5}, \mathbf{6})\mathbf{301}. \end{aligned}$$

Thus $1 = (\mathbf{2}, \mathbf{6})(\mathbf{4}, \mathbf{5})(0, 3)(5, 6)\mathbf{301}$. Hence

$$\mathbf{03} = (\mathbf{2}, \mathbf{6})(\mathbf{4}, \mathbf{5})(0, 3)(5, 6)\mathbf{11}. \quad (13)$$

We note that $1^{(\mathbf{1}, \mathbf{3}, \mathbf{0})(\mathbf{4}, \mathbf{5}, \mathbf{6})(2, 0, 6)(3, 5, 4)} = ((\mathbf{2}, \mathbf{6})(\mathbf{4}, \mathbf{5})(0, 3)(5, 6)\mathbf{301})^{(\mathbf{1}, \mathbf{3}, \mathbf{0})(\mathbf{4}, \mathbf{5}, \mathbf{6})(2, 0, 6)(3, 5, 4)}$.

So

$$1 = (\mathbf{1}, \mathbf{3}, \mathbf{0})(\mathbf{2}, \mathbf{4}, \mathbf{6})(2, 0, 5)(3, 6, 4)\mathbf{013}. \quad (14)$$

Now $\mathcal{N}^{(0)} = \mathcal{N}^0 = \langle (\mathbf{1}, \mathbf{2}, \mathbf{3}, \mathbf{6})(\mathbf{4}, \mathbf{5})(1, 2)(3, 0, 6, 5), (\mathbf{2}, \mathbf{4})(\mathbf{5}, \mathbf{6})(2, 4)(5, 6) \rangle \cong S_4$ has orbits $\{\mathbf{0}\}$, $\{1, 2, 4\}$, $\{3, 5, 6, 0\}$, and $\{\mathbf{1}, \mathbf{2}, \mathbf{3}, \mathbf{4}, \mathbf{5}, \mathbf{6}\}$. Now the 1-orbit takes the double coset $[\mathbf{0}]$ back to $[\star]$, the 4-orbit loops back to itself, since $\mathbf{00} = t\mathbf{0}$ and we need to examine the double cosets $[\mathbf{01}]$ and $[\mathbf{01}]$. Now $\mathbf{10} \sim \mathbf{01}$ (from (9)) and from (10), $[\mathbf{03}]^{(\mathbf{1}, \mathbf{3})(\mathbf{4}, \mathbf{5})(3, 5)(6, 0)} = [(\mathbf{2}, \mathbf{6})(\mathbf{4}, \mathbf{5})(0, 3)(5, 6)\mathbf{11}]^{(\mathbf{1}, \mathbf{3})(\mathbf{4}, \mathbf{5})(3, 5)(6, 0)}$ gives

$$\mathbf{01} = (\mathbf{2}, \mathbf{6})(\mathbf{4}, \mathbf{5})(0, 3)(5, 6)\mathbf{31}. \quad (15)$$

Now from (9), $((\mathbf{2}, \mathbf{6})(\mathbf{4}, \mathbf{5})(0, 3)(5, 6)\mathbf{31})^2 = (\mathbf{2}, \mathbf{5})(\mathbf{4}, \mathbf{6})(0, 3)(2, 4)$.

So $(\mathbf{2}, \mathbf{6})(\mathbf{4}, \mathbf{5})(0, 3)(5, 6)\mathbf{31}.(\mathbf{2}, \mathbf{6})(\mathbf{4}, \mathbf{5})(0, 3)(5, 6)\mathbf{31} = (\mathbf{2}, \mathbf{5})(\mathbf{4}, \mathbf{6})(0, 3)(2, 4) \implies$




Figure 4: The Cayley graph of $U_3(3)$ over $PGL_2(14)$

$$\mathbf{3131} = (\mathbf{2}, \mathbf{5})(\mathbf{4}, \mathbf{6})(\mathbf{0}, \mathbf{3})(\mathbf{2}, \mathbf{4}). \quad (16)$$

Thus $\mathbf{31} = (\mathbf{2}, \mathbf{5})(\mathbf{4}, \mathbf{6})(\mathbf{0}, \mathbf{3})(\mathbf{2}, \mathbf{4})\mathbf{13}$. Hence from (12), (14) and (15), $\mathbf{10} \sim \mathbf{01} \sim \mathbf{31} \sim \mathbf{13} \sim \mathbf{56} \sim \mathbf{65}$. Now Now $\mathcal{N}^{\mathbf{01}} = \langle y, (x * y * x^2)^2 \rangle = \langle (\mathbf{2}, \mathbf{6})(\mathbf{4}, \mathbf{5})(\mathbf{3}, \mathbf{0})(\mathbf{5}, \mathbf{6}), (\mathbf{2}, \mathbf{5})(\mathbf{4}, \mathbf{6})(\mathbf{2}, \mathbf{4})(\mathbf{3}, \mathbf{0}) \rangle$ and $\mathcal{N}t_{\mathbf{0}t_1}(\mathbf{1}, \mathbf{6})(\mathbf{2}, \mathbf{0})(\mathbf{3}, \mathbf{1})(\mathbf{4}, \mathbf{2})(\mathbf{5}, \mathbf{3})(\mathbf{6}, \mathbf{4})(\mathbf{0}, \mathbf{5}) = \mathcal{N}t_{5t_6} = \mathcal{N}t_{\mathbf{0}t_1}$. Hence $\mathcal{N}^{(\mathbf{01})} \geq \langle (\mathbf{2}, \mathbf{6})(\mathbf{4}, \mathbf{5})(\mathbf{3}, \mathbf{0})(\mathbf{5}, \mathbf{6}), (\mathbf{2}, \mathbf{5})(\mathbf{4}, \mathbf{6})(\mathbf{2}, \mathbf{4})(\mathbf{3}, \mathbf{0}), (\mathbf{1}, \mathbf{6})(\mathbf{2}, \mathbf{0})(\mathbf{3}, \mathbf{1})(\mathbf{4}, \mathbf{2})(\mathbf{5}, \mathbf{3})(\mathbf{6}, \mathbf{4})(\mathbf{0}, \mathbf{5}) \rangle \cong D_{16}$.

Now the orbits of $\mathcal{N}^{(\mathbf{01})}$ on Ω are $\{\mathbf{3}, \mathbf{1}\}$, $\{\mathbf{0}, \mathbf{1}, \mathbf{5}, \mathbf{6}\}$, and $\{\mathbf{2}, \mathbf{4}, \mathbf{5}, \mathbf{6}, \mathbf{2}, \mathbf{3}, \mathbf{4}, \mathbf{0}\}$.

Since $\mathbf{013} \sim \mathbf{1}$, $\mathbf{011} \sim \mathbf{0}$, and $\mathbf{012} = \mathbf{0.12} = \mathbf{0}\pi_{12}\mathbf{1} = \pi_{12}\mathbf{0}^{\pi_{12}}\mathbf{1} = (\mathbf{1}, \mathbf{2})(\mathbf{2}, \mathbf{0})(\mathbf{3}, \mathbf{4})(\mathbf{4}, \mathbf{6})(\mathbf{5}, \mathbf{5})(\mathbf{6}, \mathbf{3})(\mathbf{0}, \mathbf{1})\mathbf{0}^{(\mathbf{1}, \mathbf{2})(\mathbf{2}, \mathbf{0})(\mathbf{3}, \mathbf{4})(\mathbf{4}, \mathbf{6})(\mathbf{5}, \mathbf{5})(\mathbf{6}, \mathbf{3})(\mathbf{0}, \mathbf{1})}\mathbf{1} = (\mathbf{1}, \mathbf{2})(\mathbf{2}, \mathbf{0})(\mathbf{3}, \mathbf{4})(\mathbf{4}, \mathbf{6})(\mathbf{5}, \mathbf{5})(\mathbf{6}, \mathbf{3})(\mathbf{0}, \mathbf{1})\mathbf{11} \sim \mathbf{03} \in [\mathbf{01}]$, we must, therefore, have found all double cosets and the coset diagram indicated has form:

The set of all double cosets $[w] = \mathcal{N}w\mathcal{N}$, coset stabilizing subgroups $\mathcal{N}^{(w)}$, and the number of single cosets each contains are exhibited in Table 1.

Label $[w]$	Coset stabilizing subgroup $\mathcal{N}^{(w)}$	Number of cosets
$[\star]$	Since $\mathcal{N}$ is transitive on T	1
$[0]$	$\mathcal{N}^{(0)} = N^0$ $\langle y, (1, 4, 3, 5)(2, 6)(1, 4)(3, 0, 5, 6), (1, 2, 3, 6)(4, 5)(1, 2)(3, 0, 6, 5), (2, 5)(4, 6)(2, 4)(3, 0) \rangle \cong S_4$, has orbits $\{0\}, \{1, 2, 14\}, \{3, 5, 6, 0\}$ and $\{1, 2, 3, 4, 5, 6\}$ on T .	14
$[00]=[0]$	$01 \sim 10 \sim 13 \sim 31 \sim 24 \sim 42$ (Note that $[31] = [01]$)	
$[01] = [01]$ $[01] = [7, i]$	$\forall i \in \{1, 3, 0, 1, 2, 4\}$ $\forall j \in \{2, 4, 5, 6, 10, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36\}$	21
$[7, 8, j] = [7]$		

Table 3: The double cosets $[w]=\mathcal{N}w\mathcal{N}$ in $\mathcal{G}$.

6 The algorithms and programs

Algorithm I. We start with the symmetric presentation of

$$U_3(3) : 2 \cong \frac{2^{*(7+7)} : PGL_2(7)}{t = \mathbf{000}, y = (\mathbf{01})^2} \quad (17)$$

in which the three generators x, y and t correspond to the permutations a and c , and the symmetric generator t_0 , respectively. Next we compute the action of $U_3(3) : 2$ on the 36 right cosets of subgroup $PGL_2(7) = \langle x, y \rangle$ in $U_3(3) : 2$, for example, the MAGMA command **CosetAction**(U_3 , sub($U_3|x, y$)) gives J_1 in its action on 36 letters. We now form the symmetric generators $\mathcal{T} = \{t_i; i = 0, 1, \dots, 14\}$ as permutations on 36 letters and store them as ts , a sequence of length 14. Next we build cst , a sequence of length 36 whose terms are sequences of integers, representing words in the symmetric generators. These words form a complete set of coset representatives for $PGL_2(7)$ in $U_3(3)$, and correspond to the certain ordering, for example the ordering determined by the MAGMA function **CosetAction**. Represent $U_3(3) : 2 \cong \frac{2^{*(7+7)} : PGL_2(7)}{t = \mathbf{000}, y = (\mathbf{01})^2}$ as a permutation group on the right cosets of $PGL_2(7)$ in $U_3(3) : 2$ and consider a permutation p of $U_3(3) : 2$ on 36 letters. Convert p to its canonical symmetric representation form; that is, write it as a permutation of $PGL_2(7)$ on 14 letters followed by a word of length of at most 2 in the symmetric generators.

Step I $1^p = \mathcal{N} \sqrt{\quad} = \mathcal{N} \sqsupseteq \implies pw^{-1} \in \mathcal{N}$. pw^{-1} is identified with an element of $\mathcal{N}$ by its action on the fourteen cosets whose representatives are of length 1.

Step II $pw^{-1}w$ is the permutation p in its canonical symmetric representation form.

Algorithm II Convert an element x of $U_3(3)$ given in the symmetric representation form into a permutation on 36 letters. Thus x is of the form $x = nw$, where $n \in \mathcal{N}$ and w a word in the symmetric generators t_i s. Now the image of w is a permutation on 36 letters and the image of n is determined by its action on the fourteen cosets whose representatives have length 1.

Algorithm II Given two elements of $U_3(3) : 2$ symmetrically represented as (xx, uu) and (yy, vv) , the procedure **mult** uses ts and cst to return the product (zz, ww) . As described in the body of the paper the procedure **cenelt** is used to return generators for the centralizer of a given element (xx, uu) , themselves symmetrically represented. As explained above, any element of $\mathcal{G}$ can be written, not necessarily uniquely, as the product of a permutation of $\mathcal{N} \cong PGL_2(14)$ followed by a word of length at most four in the symmetric generators. In section 1.5 of [10] the first author outlines a procedure for multiplying elements represented in this fashion. It may be useful to see this process being carried out manually:

Example

$$\begin{aligned}
\pi_{00}\mathbf{12}\pi_{23}\mathbf{56} &= \pi_{00}\pi_{23}\mathbf{12}^{\pi_{23}}\mathbf{56} \\
&= \pi_{00}\pi_{23}\mathbf{0356} \\
&= (\mathbf{1, 4, 3, 2, 6, 5, 0})(1, 0, 5, 6, 2, 3, 4)\mathbf{0356} \\
&= (\mathbf{1, 4, 3, 2, 6, 5, 0})(1, 0, 5, 6, 2, 3, 4)(\mathbf{2, 4})(\mathbf{5, 6})(2, 4)(5, 6)\mathbf{3056} \\
&= (\mathbf{1, 2, 5, 0})(\mathbf{3, 4})(1, 0, 6, 4)(2, 3)\mathbf{3056} \\
&= (\mathbf{1, 2, 5, 0})(\mathbf{3, 4})(1, 0, 6, 4)(2, 3)\pi_{05}3^{\pi_{05}}\mathbf{056} \\
&= (\mathbf{1, 2, 5, 0})(\mathbf{3, 4})(1, 0, 6, 4)(2, 3)(\mathbf{1, 3})(\mathbf{2, 6})(\mathbf{3, 2})(\mathbf{4, 4})(\mathbf{5, 0})(\mathbf{6, 1})(\mathbf{0, 5})\mathbf{1056} \\
&= (\mathbf{1, 2, 5, 0})(\mathbf{3, 4})(1, 0, 6, 4)(2, 3)(\mathbf{1, 3})(\mathbf{2, 6})(\mathbf{3, 2})(\mathbf{4, 4})(\mathbf{5, 0})(\mathbf{6, 1})(\mathbf{0, 5})\mathbf{1056} \\
&= (\mathbf{1, 2, 5, 0})(\mathbf{3, 4})(1, 0, 6, 4)(2, 3)(\mathbf{1, 3})(\mathbf{2, 6})(\mathbf{3, 2})(\mathbf{4, 4})(\mathbf{5, 0})(\mathbf{6, 1})(\mathbf{0, 5})\mathbf{1056} \\
&= (\mathbf{1, 6, 4, 2})(\mathbf{2, 0})(\mathbf{3, 4, 6, 1, 5, 5, 0, 3})\mathbf{106} \\
&= (\mathbf{1, 6, 4, 2})(\mathbf{2, 0})(\mathbf{3, 4, 6, 1, 5, 5, 0, 3})(\mathbf{3, 0})(\mathbf{5, 6})(2, 6)(4, 5)\mathbf{016} \\
&= (\mathbf{1, 2})(\mathbf{2, 0})(\mathbf{3, 5})(\mathbf{4, 6})(\mathbf{5, 4})(\mathbf{6, 2})(\mathbf{0, 3})\mathbf{016} \\
&= (\mathbf{1, 2})(\mathbf{2, 0})(\mathbf{3, 5})(\mathbf{4, 6})(\mathbf{5, 4})(\mathbf{6, 2})(\mathbf{0, 3})\pi_{16}0^{\pi_{16}}\mathbf{1} \\
&= (\mathbf{1, 2})(\mathbf{2, 0})(\mathbf{3, 5})(\mathbf{4, 6})(\mathbf{5, 4})(\mathbf{6, 2})(\mathbf{0, 3})(\mathbf{1, 6})(\mathbf{2, 0})(\mathbf{3, 1})(\mathbf{4, 2})(\mathbf{5, 3})(\mathbf{6, 4})(\mathbf{0, 5})\mathbf{21},
\end{aligned}$$

which is in canonically shortest form. In this paper we have computerised such a procedure in two independent ways. The first, see Appendix A, makes full use of the ease with which MAGMA handles permutations of such low degree. Elements represented as above are transformed into permutations on 266 letters, and any group theoretic function can then be applied before transformation back into the symmetric representation. Thus, for example, the procedure *cenelt* returns generators for the centralizer of a given element symmetrically represented. One can readily write procedures to perform whatever task one chooses in MAGMA, but keep a record of the results in this short form.

The programs given in Appendix B are rather more interesting, both mathematically and computationally. To multiply two elements we first use *unify* to

express

$$\pi u . \sigma v = \pi \sigma . u^\sigma v$$

where $\pi, \sigma \in \mathcal{N}$ and u, v are words in the elements of $\mathcal{T}$, as a single sequence ss of length $11 + \text{length}(u) + \text{length}(v) \leq 19$. The first 11 entries give the permutation $\pi\sigma$ and the remainder represents a word of length ≤ 8 in the elements of $\mathcal{T}$. The procedure *canon* now puts ss into its canonically shortest form. No other representations of group elements are used; words in the symmetric generators are simply shortened by application of the relations (and their conjugates under $\mathcal{N}$). Working interactively the response is immediate.

The first type of procedure is heavily MAGMA-dependent, but can be readily modified for other packages such as GAP. On the other hand the second type of procedure, although written in MAGMA here, could have been written in any high-level language.

This work represents a first step in a programme to provide symmetric representations for interesting finite groups. Curtis has obtained suitable symmetric presentations for several sporadic simple groups; these will be computerised in a similar manner. Our next aim is ‘nested’ symmetric representations to deal with much larger examples.

Appendix

A Program 1

We start with a presentation of J_1 , based on the symmetric presentation (1), in which the three generators x , y and t correspond to the permutations a and c , and the symmetric generator t_0 , respectively. Thus the MAGMA command **CosetAction**(J , $\text{sub}(J|x, y)$) gives J_1 in its action on 266 letters. We now form the symmetric generators $\mathcal{T} = \{t_i; i = 0, 1, \dots, X\}$ as permutations on 266 letters and store them as ts , a sequence of length 11. Next we build cst , a sequence of length 266 whose terms are sequences of integers, representing words in the symmetric generators. These words form a complete set of coset representatives for $L_2(11)$ in J_1 , and correspond to the ordering determined by the MAGMA function **CosetAction**. Given two elements of J_1 symmetrically represented as (xx, uu) and (yy, vv) , the procedure **mult** uses ts and cst to return the product (zz, ww) . As described in the body of the paper the procedure **cenelt** is used to return generators for the centralizer of a given element (xx, uu) , themselves symmetrically represented.

```
U3fmt := recformat<
/* Data structure for the symmetric representation of U_3(3). */
  U3: GrpPerm,
  PGL14: GrpPerm,
  cst: SeqEnum,    // of integer sequences
  ts: SeqEnum,     // of elements of U3
  tra: SetIndx    // of elements of PGL14
>;

//-----
prodim := function(pt, Q, I)
/*
Return the image of pt under permutations Q[I] applied sequentially.
*/
  v := pt;
  for i in I do
    v := v^(Q[i]);
  end for;
```

```

        return v;
    end function;

//-----
symrep := function()
/*
Initialize the data structures for the symmetric representation of U3.
*/
G<x,y,t,s>:=Group<x,y,t,s|x^7, y^2, t^2, (x^-1 * t)^2 , (y * x)^3
, t * x^-1 * y * x \
* t * y , x^2 * y * x^3 * y * x^-4 * y * x^-4 * y*x , s^2, (s^(x^3),y),(s^(x^4\
), x*y), t * s * s^t * s, y * (s * s^(t * x^6))^2>;

    // Construct the sequence of 14 symmetric generators ts as permutations
    // on 36 letters.

    f, U3, k := CosetAction(G, sub< G | x, y,t>);
    ts := [ Id(U3): i in [1 .. 14] ];
    for i in [1..7] do
        ts[i]:=(s^(x^i))@f;
    end for;
    for i in [8..14] do
        ts[i]:=((s^t)^(x^(14-i)))@f;
    end for;

    // Construct representatives cst for the control subgroup N = PGL(2, 14)
    // as words in the symmetric generators consisting of the empty word,
    // 14 words of length one and 21 words of length two.

    S14 := SymmetricGroup(14);
    aa:=S14!(1,2,3,4,5,6,7)(14,13,12,11,10,9,8);
    bb:=S14!(2,6)(4,5)(14,10)(13,12);
    cc:=S14!(7,14)(1,8)(2,9)(3,10)(4,11)(5,12)(6,13);
    PGL14 := sub< S14 | aa, bb, cc >;

    cst := [null : i in [1 .. 36]] where null is [Integers() | ];
    for i := 1 to 14 do
        cst[prodim(1, ts, [i])] := [i];
    end for;

    tra := Transversal(PGL14, sub<PGL14 | bb,cc^(aa*bb*aa^(-1)*bb)>);
    for i := 1 to 21 do
        ss := [7,1]^tra[i];
        cst[prodim(1, ts, ss)] := ss;
    end for;

    return rec<U3fmt |
        U3 := U3, PGL14 := PGL14, cst := cst, ts := ts, tra := tra>;
end function;

//-----

//-----
sym2per := function(U3Des, x)
/*
Convert an element x of U3 in the symmetric representation into a
permutation acting on 36 letters. The image of an element of N is
determined by its action on the fourteen cosets whose
representatives have length one.
*/
    U3 := U3Des'U3; cst := U3Des'cst; ts := U3Des'ts; PGL14 := U3Des'PGL14;
    tra := U3Des'tra;

    xx := U3Des'PGL14 ! x[1]; uu := x[2];
    p := [1 : i in [1 .. 36]];

```

```

        for i := 1 to 14 do
            p[prodim(1, ts, [i])] := prodim(1, ts, [i]^xx);
        end for;
        for i := 1 to 21 do
            t := [7,1]^tra[i];
            p[prodim(1, ts, t)] := prodim(1, ts, t^xx);
        end for;
        return (U3 ! p) * &*[U3|ts[uu[j]]: j in [1 .. #uu]];
    end function;

//-----
per2sym := function(U3Des, p) /* Convert permutation p of U3 on 36
letters into its symmetric representation. The image of 1 under p
gives the coset representative for Np as a word ww in the
symmetric generators. Multiplication of p by the symmetric
generators of ww in reverse order yields a permutation which can
be identified with an element of N by its action on the 14 cosets
of length one. */
    U3 := U3Des'U3; cst := U3Des'cst; ts := U3Des'ts; PGL14 := U3Des'PGL14;
    ww := cst[1^p];
    tt := p * &*[U3|ts[ww[#ww - 1 + 1]]: 1 in [1 .. #ww]];
    zz := PGL14![rep{j: j in [1..14] | (1^ts[i])^tt eq 1^ts[j]}: i in [1..14]];
    return <zz, ww>;
end function;

//-----
cenelt := function(U3Des, x)
/*
Construct the centralizer of element x of J1 given in its symmetric representation.
An example of how all the standard procedures of MAGMA can be utilized by:
transformation to permutations, application of the procedure,
transformation back to symmetric representation.
*/
    cent := Centralizer(U3Des'U3, sym2per(U3Des, x));
    return <Order(cent), [per2sym(U3Des, c): c in Generators(cent)]>;
end function;

```

B Program 2

In this program we assume detailed knowledge of the *control subgroup* $\mathcal{N} \cong PGL_2(7)$, but use no representation of elements of J_1 other than their symmetric representation. Firstly, the procedure *unify* uses the identity $\pi u \cdot \sigma v = \pi \sigma \cdot u^\sigma v$ to combine two symmetrically represented elements (xx, uu) and (yy, vv) into a single sequence ss of length $(11 + \text{length}(uu) + \text{length}(vv))$, which represents a permutation of $\mathcal{N}$ followed by a word of length ≤ 8 in the elements of $\mathcal{T}$. The procedure *canon* then takes such a sequence and reduces it to its shortest form using the following recursive algorithm. We make use of the relations $t_i t_j t_i t_j t_i = \pi_{ij}$ and $t_i t_j t_k t_i t_j = \sigma_{ijk}$, for $\{i, j, k\}$ a *special triple* i.e. a triple fixed pointwise by an involution of $\mathcal{N}$.

The algorithm:

Step I: If two adjacent symmetric generators are equal, delete them.

Step II: If a string $t_i t_j t_i$ appears, replace it by $\pi_{ij} t_i t_j$ and move the permutation π_{ij} over the preceding symmetric generators in the standard manner.

Step III: If a string $t_i t_j t_k$ appears with $\{i, j, k\}$ a special triple replace it by $\sigma_{ijk} t_j t_i$, and move the permutation to the left as above.

Having completed the above, if $\text{length}(ss) \leq 11 + 3 = 14$, finish. Otherwise we may assume all strings $t_i t_j t_k$ have $\{i, j, k\}$ non-special.

Step IV: For each string $t_i t_j t_k t_l$ construct a permutation σ and a 5-cycle ρ such that $\sigma t_i t_j t_k$ commutes with ρ . [In fact $\sigma = \sigma_{pqr}$ where $p = i^{\pi_{jk}}, q = k^{\pi_{ij}}, r = j^{\pi_{ki}}$.] If l is not fixed by ρ we replace $t_i t_j t_k$ by $\sigma^{-1}(\sigma t_i t_j t_k) \rho^m$, where m is such that $l \in \{i, j, k\} \rho^m$. If $\text{length}(ss) \leq 15$, finish. Otherwise:

Step V: We may now assume that any string of length 5 is one of the eleven special pentads preserved by $L_2(11)$. We use the identity $267X8 = \sigma_{X64} X 26$ from [10, page 304]. i.e. for $[i, j, k, l, m]$ an even permutation of a special pentad

$$t_i t_j t_k t_l t_m = \sigma_{l j n} t_l t_i t_j,$$

where $n = l^{\pi_{ij}}$.

After each **Step** recall *canon*.

```

/*
Define the projective general
linear group PGL(2,14) as permutations of degree 14
*/
S14:=SymmetricGroup(14);
a:=S14!(1,2,3,4,5,6,7)(14,13,12,11,10,9,8);
b:=S14!(2,6)(4,5)(14,10)(13,12);
c:=S14!(7,14)(1,8)(2,9)(3,10)(4,11)(5,12)(6,13);
N:=sub<S14|a,b,c>;
/*-----
Coset Stabiliser N(0,1) /* NS71:=sub<N|N!(2, 6)(4, 5)(10, 14)(12,
13), N!(1, 13, 7, 12)(2, 9, 6, 10, 5, 11, 4, 14)(3, 8)>;
NS71:=sub<N|b,c^(a*b*a^(-1)*b)>;
/*-----
Six names for the 21-orbit
/*
ss:=[[7,1],[1,7],[13,12],[12,13],[3,8],[8,3]];
sp:=[Id(N),N!(2,5)(4,6)(14,10)(9,11),N!(7,1)(2,6,5,4)(14,9,10,11)(12,13),
N!(7,1)(2,4,5,6)(14,11,10,9)(12,13),N!(2,6)(4,5)(14,10)(12,13),
N!(2,4)(5,6)(9,11)(12,13)];
trans1:=Transversal(N,Stabiliser(N,{7,14})); prs:={@ {7,14}^x : x
in trans1 @}; pi:=N!(7,14)(1,8)(2,9)(3,10)(4,11)(5,12)(6,13);
pis:=[ pi^x : x in trans1]; trans2:=Transversal(N,NS71);

/*-----
Given two symmetrically represented elements of J1, where x and y
are permutations of N and u and v are words in the symmetric
generators, return a single sequence of length 11 + l(u) + l(v)
using the above identity.
*/
Unify := func< x, y, u, v | [ y[x[i]] : i in [1..#x] ] cat y[15..#y]
where x is Eltseq(x) cat u
where y is Eltseq(y) cat v >;
aaa:=N!(2,6)(4,5)(14,10)(13,12);
bbb:=N!(7,14)(1,8)(2,9)(3,10)(4,11)(5,12)(6,13);

```

```

Unify(aaa,bbb,u,v);
sss:=[8,9,10,11,12,13,14,1,2,3,4,5,6,7,14,7,1,8];
/*-----
Return permutation of N given by the word ijiji in the symmetric
generators. */
Pi := func< i, j | pis[Index(prs,{i,j})]>;
/*-----
For ss a sequence representing a permutation of N followed by a
word in the symmetric generators, return an equivalent sequence of
canonically shortest length.
*/
function Canon(sss)
  s:=sss;
  // Step I.
  if exists(i){ i : i in [15..#s-1] | s[i] eq s[i+1] } then
    s := Canon( s[1..i-1] cat s[i+2..#s] );
  end if;
  // Step II.
  if exists(i){ i : i in [15..#s-1] | Index(prs,{s[i],s[i+1]}) ne 0 } then
    s := Canon( [p[s[k]] : k in [1..i-1] ] cat s[i..i] cat s[i+2..#s]
      where p is Eltseq(Pi(s[i], s[i+1])) );
  end if;
  // Step III.
  if #s ge 17 then for x in trans2 do if exists(i){ i : i in [1..6] |
    [s[15],s[16]]^x eq ss[i] } then if exists(j){ j : j in [1..6] |
    [s[17]]^x eq ss[j][2] } then s:=Canon([p[s[k]] : k in [1..14]] cat
    (ss[j])^(x^(-1)) cat s[17..#s] where p is
    Eltseq((sp[i]^(-1)*sp[j])^(x^(-1))))); end if; end if; end for;end
  if;
  return s;
end function;

S14:=SymmetricGroup(14);
a:=S14!(1,2,3,4,5,6,7)(14,13,12,11,10,9,8);
b:=S14!(2,6)(4,5)(14,10)(13,12);
c:=S14!(7,14)(1,8)(2,9)(3,10)(4,11)(5,12)(6,13);
N:=sub<S14|a,b,c>; NS71:=sub<N|N!(2, 6)(4, 5)(10, 14)(12, 13),
N!(1, 13, 7, 12)(2, 9, 6, 10, 5,\
  11, 4, 14)(3, 8)>;
ss:=[[7,1],[1,7],[13,12],[12,13],[3,8],[8,3]];
sp:=[Id(N),N!(2,5)(4,6)(14,10)(9,11),N!(7,1)(2,6,5,4)(14,9,10,11)(12,13),
N!(7,1)(2,4,5,6)(14,11,10,9)(12,13),N!(2,6)(4,5)(14,10)(12,13),
N!(2,4)(5,6)(9,11)(12,13)];
trans1:=Transversal(N,Stabiliser(N,{7,14})); prs:={@ {7,14}^x : x
in trans1 @}; pi:=N!(7,14)(1,8)(2,9)(3,10)(4,11)(5,12)(6,13);
pis=[ pi^x : x in trans1]; Pi := func< i, j |
pis[Index(prs,{i,j})]>; trans2:=Transversal(N,NS71); function
Canon(sss)
  s:=sss;
  // Step I.
  if exists(i){ i : i in [15..#s-1] | s[i] eq s[i+1] } then
    s := Canon( s[1..i-1] cat s[i+2..#s] );
  end if;
  // Step II.
  if exists(i){ i : i in [15..#s-1] | Index(prs,{s[i],s[i+1]}) ne 0 } then
    s := Canon( [ p[s[k]] : k in [1..i-1] ] cat s[i..i] cat s[i+2..#s]
      where p is Eltseq(Pi(s[i], s[i+1])) );
  end if;
  // Step III.
  if #s ge 17 then if exists(i,j,x){<i,j,x> : i in [1..6], j in
  [1..6] , x in trans2 | ss[i]^x eq [s[15],s[16]] and ss[j][2]^x eq
  s[17]} then s:=Canon([p[s[k]] : k in [1..14]] cat [ss[j][1]^x
  cat s[18..#s] where p is Eltseq((sp[i]^(-1)*sp[j])^x)); else if
  exists(i,j,x){<i,j,x> : i in [1..6], j in [1..6] , x in trans2 |
  ss[i]^x eq [s[15],s[16]] and Index(prs,{ss[j][2]^x,s[17]}) ne 0}
  then s:=Canon([p[s[k]] : k in [1..14]] cat
  [ss[j][1]^x(Pi(ss[j][2]^x,s[17]))] cat [ss[j][2]^x cat s[18..#s]
  where p is Eltseq((sp[i]^(-1)*sp[j])^x*Pi(ss[j][2]^x,s[17]))); end

```

```

if; end if; end if;
return s;
end function;
/*-----
Return the product of two symmetrically represented elements of J1
*/ Prod := function( x, u, y, v)
    t := Canon(Unify(x, u, y, v));
    return N!t[1..14], t[15..#t];
end function;

/*-----
Return the inverse of a symmetrically represented element of J1
*/
Invert := func< x, u | x^-1, [u[#u-i+1]^(x^-1) : i in [1..#u]] >;

```

References

- [1] J. N. Bray, R. T. Curtis, 'Double coset enumeration of symmetrically generated groups', *J. Group Theory* 7 (2004) 167-185.
- [2] John Cannon and Catherine Playoust, 'An Introduction to MAGMA', *School of Mathematics and Statistics, University of Sydney*. (1993).
- [3] J.H.Conway, 'Three lectures on exceptional groups', M.Powell and G.Higman Eds. *Finite Simple Groups*, *Academic Press*,(1970), pp.215–247.
- [4] J.H.Conway, R.T.Curtis, S.P.Norton, R.A.Parker and R.A.Wilson, *An ATLAS of Finite Groups*, *Oxford UP* (1985).
- [5] R.T.Curtis, 'Symmetric Presentations I: Introduction, with particular reference to the Mathieu groups M_{12} and M_{24} ,' *Proceedings of the L.M.S. Durham Conference on 'Groups and Combinatorics,'* (1990).
- [6] R. T. Curtis. Symmetric generation of Higman-Sims group. *J. Algebra* 171 (1995), 567-586.
- [7] R. T. Curtis. Symmetric generation and existence of the Janko group J_1 . *J. Group Theory* 2 (1999), 355-366.
- [8] R. T. Curtis and Z. Hasan, 'Symmetric representation of the elements of the Janko group J_1 ', *J. Symbolic Computation* (1996) 22, 201-214.
- [9] Abdul Jabbar, 'Symmetric presentations of the subgroups of the Conway groups and related topics, Ph.D thesis, University of Birmingham, England (1992).
- [10] R.T.Curtis, 'Symmetric Presentations II: The Janko group J_1 ', *J. London Math. Soc.* (1993).
- [11] Z.Janko, 'A new finite simple group with Abelian Sylow 2-subgroups,' *Proc. Nat. Acad. Sci. USA*, 53(1965), pp. 657–658.
- [12] Z.Janko, 'A new finite simple group with Abelian Sylow 2-subgroups, and its characterisation,' *J. Algebra* 3 (1966) 147–186.
- [13] D.Livingstone, 'On a permutation representation of the Janko group', *J. Algebra*, 6 (1967) 43–55.
- [14] E.Mathieu, 'Memoire sur l'etude des fonctions de plusieurs quantites,' *J. Math. Pures Appl.* 6 (1861) 241–243.
- [15] E.Mathieu, 'Sur la fonction cinq fois transitive de 24 quantites,' *J. Math. Pures Appl.* 18 (1873) 25-46.
- [16] R. A. Parker, 'The Computer Calculation of Modular Characters (The MEATAXE),' *Computational Group Theory*, Ed. Michael D. Atkinson, *Academic Press*(1984).

- [17] R.A.Wilson, ‘An Atlas of sporadic group representations,’ *Proceedings of the Birmingham conference ‘The ATLAS ten years on,’* Eds. R. T. Curtis and R. A. Wilson (pending).